

PREVENT ANAEMIA **IN** PREGNANCY




DENA
AITO EJE
NINU
OYUN

BLOOD LEVEL



> 11G/L or > 33%


IWON EJE



< 7G/L or < 21%

SOME CAUSES OF ANAEMIA IN PREGNANCY ARA AWON OHUN TI O NFA AITO EJE NINU OYUN

SHORT PREGNANCY INTERVAL (AIFI ETO SI OMO BIBI)





MALARIA (AISAN IBA)



WORM INFESTATION (ARAN INU)



(Unwashed Fruit)
Eso ti ako fo



PREVENTION

NUTRITIOUS MEAL (OUNJE TI O NSE ARA LOORE)



THE GOOD AND THE BAD MEAL (OUNJE TODARA ATI EYI TI KODARA)



NOODLES/PASTA
(NUDULU /PASITA)



JUNK
(WEWE)



DRINKS
(OTI ELERIN-DODO)



FRUITS
(AWON ESO)




VEGETABLES
(AWON EWEBE)



PROTEIN & IRON
(AFUNNI LEJE)



HAEMETINICS DRUG/SUPPLEMENT INTAKE (OGUN TI ONDENA AITO EJE)



PROPHYLACTIC ANTIMALARIA (IPT_p-SP)(OOGUN IBA)



PROPHYLACTIC IPTP-SP (OGUN TI O NDENA AISAN IBA NINU OYUN)

**THIS DRUG ARE GIVEN AT (4-9) MONTHS IN PREGNANCY
TO PREVENT MALARIA**



PREVENTION

FAMILY PLANNING

(IFETO SOMO BIBI)



WORM PREVENTION


DIDENA AARAN



CLEAN ENVIRONMENT (IMOTOTO AYIKA)



DIRTY ENVIRONMENT (AYIKA TI ODOTI)



CONSEQUENCIES OF ANAEMIA IN PREGNANCY ATUBOTAN AITO EJE NINU OYUN



ANAEMIC BABY (OMO TI EJE RE KO PE OJU OSUWON)



CONSEQUENCIES OF ANAEMIA IN PREGNANCY

ATUBOTAN AITO EJE NINU OYUN



ANAEMIC/SICK MOTHER
(IYA TI EJE RE KO GBE OSUWON/TI O NSE AISAN)



PREMATURE BABY
(OMO TI KOGBO)



DEAD MOTHER
(OKU IYA)



MALFORMED BABY
(OMO TI EYA ARA WON KOPE)



DEAD BABY
(OKU OMO)